



Oral potassium supplementation in surgical patients

Alison J. Hainsworth*, Piers A. Gatenby

General Surgical Unit, St Mary's Hospital, Imperial College NHS Trust, UK

ARTICLE INFO

Article history:

Received 4 March 2008

Accepted 18 March 2008

Published online 28 March 2008

Keywords:

Hypokalaemia

Dietary modification

ABSTRACT

Hospital inpatients are frequently hypokalaemic. Low plasma potassium levels may cause life threatening complications, such as cardiac arrhythmias. Potassium supplementation may be administered parenterally or enterally. Oral potassium supplements have been associated with oesophageal ulceration, strictures and gastritis. An alternative to potassium salt tablets or solution is dietary modification with potassium rich food stuffs, which has been proven to be a safe and effective method for potassium supplementation. The potassium content of one medium banana is equivalent to a 12 mmol potassium salt tablet. Potassium supplementation by dietary modification has been shown to be equally efficacious to oral potassium salt supplementation and is preferred by the majority of patients. Subsequently, it is our practice to replace potassium using dietary modification, particularly in surgical patients having undergone oesophagogastricectomy or in those with peptic ulcer disease.

© 2008 Surgical Associates Ltd. Published by Elsevier Ltd. All rights reserved.

Hospital inpatients are frequently hypokalaemic.¹ Hypokalaemia is more common in patients with increased potassium losses via the kidney, for example due to diuretics² or tubular dysfunction, or via the gastrointestinal tract, secondary to diarrhoea, vomiting or bowel preparation.^{3,4} Potassium supplementation is often given to correct hypokalaemia and may be administered parenterally or enterally (via gastric or jejunal feeding tubes or taken orally).

Oral potassium supplements have been associated with oesophageal ulceration.^{5–7} They have also been implicated in causing oesophageal strictures^{5,8} and gastritis.⁹

Maintenance potassium requirements are 1 mmol/kg/day and when plasma potassium levels are low or falling the minimal excretion of ionized potassium is 0.2 mmol/kg/day.^{1,10} Low plasma potassium levels can cause life threatening cardiac arrhythmias and weakness of respiratory muscles.¹⁰ All patients with potassium levels below the lower limit of normal should be treated. In some patients (for example, those with cardiac disease, atrial fibrillation or those on digoxin) it is prudent to maintain a potassium level towards the upper limit of normal.¹

Dietary modification involving potassium-rich food-stuffs has long been recognized as a potential method for potassium supplementation.¹¹ Norris et al. proved this as a safe, cheaper and equally efficacious alternative to potassium salt medication and the majority (79%) of patients preferred diet supplementation.¹²

Oral potassium tablets typically contain around 12 mmol of potassium and should be administered with food to reduce their alimentary side effects, which include nausea and vomiting.¹² This is equivalent to one medium banana.¹²

Bananas are a well accepted source of potassium; 1 g of banana contains 1 mmol of potassium.¹³ Other potassium-rich foods include baked potato (one baked potato contains 22 mmol potassium) and prune juice (9 mmol in 150 ml) or orange juice (4 mmol in 150 ml).¹²

The potassium deficit for a reduction in serum potassium concentration is variable, but is typically around 100 mmol potassium for a 0.3 mmol/l reduction in serum potassium.¹ Patients with hypokalaemia typically receive approximately 72 mmol of potassium salt supplementation a day in the form of six dissolvable tablets. This is alternative to six

* Corresponding author.

E-mail address: alison.hainsworth@ic.ac.uk (A.J. Hainsworth).

1743-9191/\$ – see front matter © 2008 Surgical Associates Ltd. Published by Elsevier Ltd. All rights reserved.

doi:10.1016/j.ijsu.2008.03.007

bananas. Sando-K contains 12 mmol potassium and costs £1.53 for 20 tablets,¹⁴ a cost of 0.6 pence/mmol potassium. Bananas in my local supermarket cost 77 p/kg (6 medium bananas) equivalent to 1.1 pence/mmol potassium. A 20 mmol ampoule of potassium chloride costs 48 pence,¹⁴ the equivalent of 2.4 pence/mmol.

Subsequently, our practice has been to give oral potassium supplementation by dietary modification to surgical patients when this is possible, particularly in those patients who have undergone oesophagogastric resection and those with upper alimentary disease such as reflux oesophagitis and peptic ulcer disease.

Conflict of interest

None declared.

Funding

None.

Ethical approval

Not required.

REFERENCES

- Gennari F. Hypokalaemia. *N Engl J Med* 1998;**339**(7):451–8.
- Arif M, Eaton S. The effect of oral potassium supplements on plasma potassium in normokalaemic and hypokalaemic patients. *J Pak Med Assoc* 1977;**27**(11):424–6.
- Downing T, Bennion R, Sadeghi A. Effect of preoperative colon preparation on serum potassium. *Am Surg* 1983;**49**(8):414–6.
- Ritsema G, Eilers G. Potassium supplements prevent serious hypokalaemia in colon cleansing. *Clin Radiol* 1994;**49**(12):874–6.
- Pemberton J. Oesophageal obstruction and ulceration caused by oral potassium therapy. *Br Heart J* 1970;**32**(2):267–8.
- Rosenthal T, Adar R, Militanu J, Deutsch V. Esophageal ulceration and oral potassium chloride ingestion. *Chest* 1974;**65**:463–5.
- Stanley A, Eade O, Hardwick D. Oesophageal ulceration secondary to potassium tablets. *Scott Med J* 1994;**39**(4):118–9.
- Lambert J, Newman A. Ulceration and stricture of the esophagus due to oral potassium chloride (slow release tablet) therapy. *Am J Gastroenterol* 1980;**73**(6):508–11.
- McLoughlin J. Effects on upper gastrointestinal mucosa of three delivery systems of potassium as supplement to frusemide administration. *J R Soc Med* 1985;**79**(6):459–62.
- Groeneveld J, Sikpkens Y, Lin S, Davids M, Halperin M. An approach to the patient with severe hypokalaemia: the potassium quiz. *QJM* 2005;**98**(4):305–16.
- Saggar-Malik A, Cappuccio F. Potassium supplements and potassium-sparing diuretics. A review and guide to appropriate use. *Drugs* 1993;**46**(6):986–1008.
- Norris W, Kunzelman K, Bussell S, Rohweder L, Cochran R. Potassium supplementation, diet vs pills. *Chest* 2004;**125**:404–9.
- Clements M, Levine M, Black R, Hughes T, Rust J, Tome F. Potassium supplements for oral diarrhoea regimens. *Lancet* 1980;**2**(8199):854.
- London, editor. *British National Formulary*. 54 ed. BMJ Publishing Group; 2007.